

### AP Experiments Liaison

- Background Signals
- CDEV, BERT
- ZDC
- Cross Sections
- Vertex

RHIC Retreat 2005 Angelika Drees


# Background contamination with collision rates during Cu-run





- top: STAR blue halo and yellow halo signals, bottom: PHENIX scintillator signals (N5&6, S5&6)
- rates drop to less than 20% during an uncogging experiment in PHENIX and STAR blue halo (yellow halo?), N6 was somewhat different
- for the remainder of the run we successfully scaled the PHENIX backgrounds with the colllision rate to compensate for this:
  - $bkgd = bkgd-scalef \times coll$
- scalef = 0.17 to 0.28


### STAR pp background signal quality




- uncogging beams with fill patterns > 56 bunches leaves some remaining collisions
- blue and yellow backgrounds drop to almost zero
- amount of collimation contamination is very different in blue and yellow signal: 1/9 (blue) and about 1/3 (yellow)
- the blue background signal is dominated by collisions!

# STAR Background Signals during transverse steering (vernier scan)


- uncogging experiment can be confirmed with transverse steering experiments (aka vernier scans;))
- top: scan early in the pp run, bottom: scan later in the pp run
- while yellow even increases for very missteered beams, blue behaves almost like the collision rate (BBC, red).
- blue Halo does not give a good background measure! New blue background is clearly not contaminated by collisions







## STAR backgrounds "new" vs. "old"

- good or bad conditions are based on the 'old' blue Halo and yellow Halo signals: BBC/(BH+YH) > 10.
- new signals are timed into the abort gap of the other ring and scaled to the total number of bunches
- both yellow signals are close but blue signals were very different, not clear what to do
- needs STAR expert to look into and to study




## STAR backgrounds "new" vs. "old"






- □ left: store with 63x63 bunches, right: store with 56x56 bunches (205 GeV!)
- Yellow is more or less consistent in both cases, blue is clearly very different
- however, I've seen some 56x56 stores with larger differences in the two blue background signals than in the two 205 GeV stores.

# Reliability of STAR backgrounds for steering and collimation






- top: ZDC collision rate from BRAHMS, PHENIX and STAR, STAR is not optimized
- when optimization is attempted (using yellow beam), ZDC rates are clearly overshooting the goal
- during that attempt there is no increase in the yellow background!
- could that be due to saturation? (I've seen even higher rates though)
- final optimization only possible after collimation (and blue beam was used => some spike during steering!)

## Summary Backgrounds

- PHENIX scintillator backgrounds are reliable, useful and consistent for steering and collimation
- STAR backgrounds are varying and at times dominated by collision signals
- we lack support from some expert to study and understand those signals (fill pattern dependence?)
- collimation and background reduction at STAR is difficult (if not impossible) under those conditions
- there were no background issues with any of the other experiments (except, maybe, pressure rise caused backgrounds) due to large beta\* values


## CDEV: Getting data online to experiments

- list of CDEV parameters
  was cleaned up before
  the beginning of this run
- there are PET pages available to experiment experts to check on all CDEV devices & parameters online (common and individual)
- there were only minor issues with CDEV this year (correct me if I'm wrong!)




## Online communications with Exp.

- BERT is routinely used by MCR and experiments
- phone contacts are still important and frequent though
- does this tool need improvements or changes? Let us know ...




#### ZDC collision rates

- steering, optimization (even collimation) and bookkeeping is based on the ZDC collision rate
- we check the goal by the ratio of the various experiments according to their beta\* values
- during the 205 GeV run ZDC rates clearly do not scale with beta\* (10/2)!
- unclear why, are STAR & PHENIX too high? PHOBOS to low?
- need input from all experiments to find out: difficult to get
- ZDC readout should be a C-AD system, HV or thresholds change with energy and species



#### ZDC readout as C-AD device


- a C-AD copy of the existing readout electronics is already in place since the beginning of the Cu run
- we have no way (yet) to determine the right threshold => work in progress (see T. Russo talk tomorrow)
- □ signal evolution is consistent with collision signal ©
- □ signal is smaller and noisier than experimenter signal (not understood yet) ⊗
- we are working on timing with the beam synchronous clock for background reduction



#### Cross Section Measurements

- cross section measurements needed for bookkeeping (and experimental purposes)
- measured by (more or less) regular vernier scans
- cross section measurement at 205 GeV yields about factor 4 higher than 100 GeV: approx. 1.6 mb (vs. 0.39 mb), not yet understood (work in progress with PHENIX)
- pp cross section in ZDCs changed from 0.33 mb to 0.39 mb this year, not understood
- open issue from pp 2004 of decreasing measured luminosity with increase of delivered luminosity, so far no input from experiments (B. Surrow?)







bump amplitude (mm)

horizontal scan @ 205 GeV


#### Mean Vertex Evolution






- □ top: early April, bottom: Jun 01-09
- we requested online zvertex information from all experiments before the run
- we got at some point zand x,y-vertex from PHOBOS, z vertex from PHENIX, no data from STAR and BRAHMS


#### Z-vertex RMS variations






- same time periods as before (data from PHENIX)
- larger variations early in the run than later
- RMS values (from PHENIX online fits!) are typically around 21 cm

## Z-vertex histograms



- 'raw' data from vertex distributions are available from PHOBOS and PHENIX
- top: 1<sup>st</sup> vertex in store, bottom: last vertex in store
- fits do not give the exact same results as PHENIX online fits, varies by up to 20% => need to talk to someone
- vertex distribution is constant to 1<sup>st</sup> order


#### zoom into one store




- though the RMS width is relatively stable there is some slight increase during the store
- increase rate is about 1mm/hour (not significant)
- data is from PHENIX online fits

#### Transverse vertex @ store

- PHOBOS delivered transverse vertex distributions
- so far we looked into PHOBOS online fit data only
- top: vertical vertex position from PHOBOS, bottom: BPM readings
- □ from vertex: -1.35 mm
- □ from BPM: -0.97 mm (blue) and -1.15 mm (yel)
- consistent within 0.4 mm (!)
  (similar in horizontal plane and other stores) ©
- unfortunately, PHOBOS is the only one to deliver this data and here the DX BPMs are the most consistent with each other!





### Summary

- Backgrounds with PHENIX: no issues
- Backgrounds with STAR: confusing, lacks support
- CDEV: no issues this year (?)
- ZDCs: HV and/or thresholds keep being an issue, transformation into C-AD system in progress -> how to change thresholds?
- Cross sections: needed for bookkeeping, open issues from pp\_04
- collaboration on Xsec with PHENIX is going on ©
- no collaboration with STAR established (yet?)
- vertex data from experiments:
  - no response from BRAHMS ⊗
  - z-vertex data from PHENIX ©
  - x,y,z vertex data from PHOBOS during Cu run ©
  - no data from STAR (though x,y,z were promised) ⊗